

# PRV

PATENT- OCH REGISTRERINGSVERKET  
Patentavdelningen

SE00/A67

PCT/SE 00/01767<sup>712</sup>  
10/070412

REC'D 01 NOV 2000

WIPO

PCT



## Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande AstaCarotene AB, Gustavsberg SE  
Applicant (s)

(21) Patentansökningsnummer 9903336-7  
Patent application number

(86) Ingivningsdatum 1999-09-17  
Date of filing

Stockholm, 2000-10-20

För Patent- och registreringsverket  
For the Patent- and Registration Office

Anita Södervall  
Anita Södervall

Avgift  
Fee

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

29295/BN

# **DNA construct and its use.**

The present invention relates to a new DNA construct for transformation into oilseed plants. The DNA construct comprises nucleotide sequences encoding peptides with enzyme activities necessary for the high-level production and esterification of keto group-containing xanthophylls in oilseed plants.

## **Background of the invention**

Carotenoids are produced *de novo* by plants, fungi, algae and some bacteria. A number of biosynthetic steps are needed for the biological production of the carotenoids. There are two chemically different groups of carotenoids, namely carotenes containing only carbon and hydrogen molecules and xanthophylls containing oxygen in the molecule in addition to carbon and hydrogen.

The xanthophylls, and particularly astaxanthin (3,3'-dihydroxy- $\beta$ - $\beta$ -carotene-4,4'-dione), are often colored pigments and are used as such or as anti-oxidants.

Carotenes are biological precursors for the production of the oxygen-containing xanthophylls. There are two types of enzymes responsible for the introduction of hydroxy groups and keto groups into the carotenes, namely hydroxylases and ketolases, respectively.

The keto group-containing xanthophyll astaxanthin, which has keto and hydroxy groups, is biosynthetically produced from beta-carotene.

Large-scale production of xanthophylls from natural sources is at present performed by AstaCarotene AB, Gustavsberg, Sweden, by cultivation of the alga *Haematococcus pluvialis* for the production of astaxanthin in esterified form.

It would be desirable to be able to produce keto group-containing xanthophylls particularly astaxanthin, in oilseed plants. Oilseed plants have naturally  $\beta$ -carotene hydroxylases but lack  $\beta$ -carotene C-4-oxygenase enzymes or ketolases.

## **Description of the invention**

The present invention provides DNA constructs enabling and promoting production of keto group containing xanthophylls, especially astaxanthin, in oilseed plants, such as rape, sunflower, soybean and mustard. The DNA construct is transformed into the oilseed plant cell for expression of a protein or fused protein which has an enzyme activity enabling keto group insertion into a carotene or hydroxy carotene for the biosynthetic production of a keto group containing xanthophyll, such as cantaxanthin ( $\beta$ , $\beta$ -carotene-4,4'-dione) and/or astaxanthin. Use is thus made of the biosynthetic pathway of the oilseed plant to

produce carotenoids. The naturally occurring synthesis of carotenoids involves a number of enzymes, namely 1-D-deoxyxylulose 5-phosphate synthase, isopentenyl pyrophosphate:dimethylallyl pyrophosphate isomerase, geranylgeranyl pyrophosphate synthase, phytoene synthase, phytoene desaturase, zeta-carotene desaturase, lycopene beta-cyclase,  $\beta$ -carotene hydroxylase, and  $\beta$ -carotene C-4-oxygenase. Genes coding for peptides having these enzymatic activities may be inserted into the DNA construct of the invention, one or several per construct, to promote high-level production in the transgenic oilseed plant. In case only one enzyme coding gene is inserted per plant, two or more plants may be sexually interbred to produce plants containing all the desired enzyme activities.

Thus, the present invention is directed to a DNA construct comprising in the 5' to 3' direction of transcription operably linked a promoter region directing transcription to the seed of an oilseed plant, a nucleotide sequence coding for at least one peptide with enzyme activity necessary for keto group containing xanthophyll production and esterification in an oilseed plant and a transcriptional termination region.

In a preferred embodiment of the invention the DNA construct additionally comprises between the promoter region and the nucleotide sequence coding for at least one peptide with enzyme activity a nucleotide sequence coding for a transit peptide directing the translated fusion polypeptide to the chloroplast of the oilseed plant.

The DNA construct is preferably such that the promoter is a napin promoter, the peptide with enzyme activity necessary for keto group containing xanthophyll production is selected from the group consisting of peptides with 1-D-deoxyxylulose 5-phosphate synthase, isopentenyl pyrophosphate:dimethylallyl pyrophosphate isomerase, geranylgeranyl pyrophosphate synthase, phytoene synthase, phytoene desaturase, zeta-carotene desaturase, lycopene beta-cyclase,  $\beta$ -carotene hydroxylase, and  $\beta$ -carotene C-4-oxygenase activity. To promote esterification of astaxanthin a nucleotide sequence coding for a peptide with acyl transferase activity may be included in the group.

In a preferred embodiment of the DNA construct according to the invention the nucleotide sequence coding for a peptide with enzyme activity is a nucleotide sequence coding for a N-terminally truncated  $\beta$ -carotene C-4-oxygenase gene from the alga *Haematococcus pluvialis*.

An example of the DNA construct of the invention is presented in the sequence listing as SEQ ID NO:1 and in Fig.1.

The present invention is also directed to a transgenic oilseed plant cell comprising the DNA construct of the invention, and preferably the oilseed plant is selected from the group consisting of rape, sunflower, soybean and mustard.

The invention is additionally directed to transgenic oilseed plant-produced xanthophyll, e.g. canthaxanthin and astaxanthin.

A preferred aspect of the invention is directed to transgenic oilseed plant-produced astaxanthin esters.

The present invention will now be illustrated with reference to the DNA construct disclosed in the sequence listing and in Fig.1, and the following description of embodiments. However, the invention is not limited to these exemplifications.

#### Short description of the drawings

Fig.1 illustrates the nucleotide sequence of the DNA construct comprising the napin promoter, the chloroplast localization signal, the N-terminally truncated  $\beta$ -carotene C-4-oxygenase gene and the termination sequence, and the deduced amino acid sequences of the transit peptide and the  $\beta$ -carotene C-4-oxygenase.

#### Description of embodiments

The invention is illustrated by production of astaxanthin in the seed of oilseed rape. The astaxanthin produced in the seed of the transgenic plant is extracted as part of the extracted oil. By use of conventionally used protocols for *Agrobacterium tumefaciens* mediated transformation such as described by (Hoekema et al.1983, An et al. 1986, Fry et al. 1987, DeBlock et al. 1988, Radke et al.1988, or Moloney et al. 1989) transgenic plants are produced having a chimeric DNA construct that is genetically inherited and is able to produce astaxanthin. The nucleotide sequence of the chimeric DNA construct consist of four parts of different genetic origin namely: (1) a promoter, (2) a localization signal, (3) a  $\beta$ -carotene C-4-oxygenase coding region and (4) a termination sequence.

The napin promoter directs transcription to the seed of oilseed rape (Stålberg et al 1996). This promoter was coupled to a localization signal similar but not identical to a transit peptide (TP) of Rbcs1a (Krebbers, 1988) that directs the translated product of a fused gene to the chloroplast. The promoter and the TP sequence were ligated to a part of the coding sequence of a ketolase gene BCK (Kajiwara et al. 1995). This enzyme oxygenates  $\beta$ -carotene to canthaxanthin, (Fraser et al. 1997). The chimeric DNA construct was then coupled to a suitable termination sequence, e.g. that of the *Agrobacterium tumefaciens* nopaline synthase gene (the nos 3' end)(Bevan et al. 1983), as illustrated in Fig.1.

### Cellular storage of Astaxanthin

The storage of large amounts of free astaxanthin in plants will be difficult due to toxic effects of the molecule as it intercalates in the plant membranes. An effective esterification of astaxanthin to fatty acids enables storage of the esterified molecules in triacylglycerol containing oleosomes. Thus, an acyl transferase can be claimed to be of  
 5 fundamental importance for the process, as is proteins that can mediate transport of different forms of astaxanthin from the chloroplast to the vesicles.

### Sequences and oligonucleotides used in the construction of the DNA construct

#### 1. *Napin promoter (GeneBank ACCESSION No. J02798)*

10 This promoter sequence, a 1145 base pair fragment including the 5' leader sequence has a unique HindIII site at the 5' end. The 3' end was synthesized with an additionally 6 nucleotide BamHI site.

#### 2. *Transit peptide similar to RBCS1a (GeneBank ACCESSION No. X13611, X14565)*

The transit peptide (TP) was amplified by PCR from -28 to the end of the transit  
 15 cleavage aa=54/55 site of the Rbcs1a gene. The 5' end was synthesized with a BamHI site and similarly the 3' sequence was synthesized with a XbaI site. The two following oligonucleotides were used for the PCR amplification.

#### BamHI

20 5' primer: TP1 5'AGAC GGATCC TCAGTCACACAAAGAGTA 3'

#### SacI XbaI

3' primer: TP2 5'GTTC GAGCTC TCTAGA CATGCAGTTAACGC 3'

#### 3. *BCK ( $\beta$ -carotene C-4 oxygenase) (Genebank ACCESSION No. D45881)*

25 The BCK fragment was amplified by PCR including a 5' XbaI site and was ligated to the TP already described. The 5' primer (BCK1) used for PCR, is homologous to the BCK sequence from nucleotide 264 and the 3' oligonucleotide (Ax40) ends with a stop codon and was synthesized with a SacI restriction site for cloning. The synthesized fragment was fused to the TP as shown in Fig 1.

30 Oligonucleotides used for PCR:

#### XbaI

5' primer: BCK1 5'ACAG TCTAGA ATGCCATCCGAGTCGTCA 3'

#### SacI

3'primer: AX40 5'CACCGAGCTCCATGACACTCTTGTGCAGA 3'

# Description of SEQ ID NO:1 and SEQ ID NO:2

The sequences shown in Fig.1 are the same as the two sequences which are shown in the sequence listing.

The SEQ ID NO:1 is a nucleotide sequence composed of the following features:

5		Nucleotide No.
	Cloning site HindIII	1-6
	Napin Promoter	1-1145
	Cloning site BamHI	1146-1151
	Transit peptide leader	1152-1178
10	Transit peptide coding	1179-1347
	Cloning site XbaI	1348-1353
	$\beta$ -carotene C-4-oxygenase	1354-2217
	$\beta$ -carotene C-4-oxygenase 3' untranslated	2218-2266
	Cloning site SacI	2267-2272
15	Nopaline synthetase termination	2273-2536
	Cloning site EcoRI	2538-2543

The SEQ ID NO: 2 is a deduced amino acid sequence of the fusion protein of the transit peptide and the peptide with  $\beta$ -carotene C-4-oxygenase activity.

## References

- An G, Watson BD, Chiang CC (1986), Transformation of tobacco, tomato, potato and  
5 *Arabidopsis-thaliana* using a binary vector system. *Plant Physiology* 81 (1) 301-305.
- Bevan M, Barnes WM and Chilton MD (1983). Structure and transcription of the nopaline  
synthase gene region of T-DNA. *Nucleic Acids Res.* 11 (2), 369-385 .
- 10 DeBlock M, DeBrouwer D, Tenning P (1989). Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the BAR and NEO genes in transgenic plants *Plant Physiology* 91:2, 694-701.
- Fraser PD, Miura Y, Misawa N, (1997). In vitro characterization of astaxanthin biosynthetic  
15 enzymes. *J Biol Chem.* Mar 7;272(10):6128-35.
- Fry J, Barnason A, and Horsch RB, (1987). Transformation of *Brassica napus* with *Agrobacterium tumefaciens* based vectors. *Plant Cell Reports* 6:321-325.
- 20 Hoekema A, Hirsch PR, Hooykas PJJ Schilperoort, (1983). A binary vector strategy based on separation of vir and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. *Nature* vol 303, 179-180.
- Josefsson LG, Lenman M, Ericson ML and Rask L, (1987). Structure of a gene encoding the  
25 1.7 S storage protein, napin, from *Brassica napus*. *J. Biol. Chem.* 262 (25), 12196-12201.
- Kajiwara S, Kakizono T, Saito T, Kondo K, Ohtani T, Nishio N, Nagai S and Misawa N.  
(1995). Isolation and functional identification of a novel cDNA for astaxanthin biosynthesis  
from *Haematococcus pluvialis*, and astaxanthin synthesis in *Escherichia coli* *Plant Mol. Biol.*  
30 29 (2), 343-352.

Krebbers E, Seurinck J, Herdies L, Cashmore AR and Timko MP, (1988). Four genes in two diverged subfamilies encode the rubulose-1, 5-bisphosphate carboxylase small subunit polypeptides of *Arabidopsis thaliana* Plant Mol. Biol. 11, 745-759.

- 5 Moloney M, Walker JM and Sharma KK, (1989). High efficiency transformation of *Brassica napus* using *Agrobacterium* vectors. Plant Cell Reports 8:238-242.

- 10 Radke SE, Andrews BM, Moloney MM, Crouch ML, Kridl JC, Knauf VC (1988), Transformation of *Brassica napus* using *Agrobacterium tumefaciens* – Developmentally regulated Expression of a reintroduced napin gene. TAG, 75: (5) 685-694 .

Pua E-C, Mehra-Palta A, Nagy F and Chua N-H, (1987). Transgenic plants of *Brassica napus*. Biotechnology vol 5, 815-817.

- 15 Stålberg K, Ellerstöm M, Ezcurra I, Ablov S, Rask L (1996). Disruption of an overlapping E-box/ABRE motif abolished high transcription of the napA storage-protein promoter in transgenic *Brassica napus* seeds. Planta 199(4):515-9.



## SEQUENCE LISTING

<110> AstaCarotene AB

<120> DNA construct and its use

<130> 29295-AstaCarotene

<140>

<141>

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 2543

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: napin promoter  
+ chloroplast localization signal + beta-carotene C-4 oxygenase  
coding sequence + termination sequence

<220>

<221> promoter

<222> (1)..(1145)

<220>

<221> transit\_peptide

<222> (1179)..(1347)

<220>

<221> CDS

<222> (1179)..(2217)

<220>

<221> terminator

<222> (2273)..(2536)

<400> 1

aagctttctt catcggtgat tgattccttt aaagacttat gtttcttattc ttgctttctga 60  
ggcaagtatt cagttaccag ttaccactta tattctggac tttctgactg catcctcatt 120  
tttccaacat tttaaatttc actattggct gaatgcttct tctttgagga agaaacaatt 180  
cagatggcag aaatgtatca accaatgcat atatacaaat gtacctcttg ttctcaaaac 240  
atctatcgga tggttccatt tgctttgtca tccaattagt gactacttta tattattcac 300  
tcctctttat tactattttc atgcgagggt gccatgtaca ttatatttgt aaggattgac 360  
gctattgagc gtttttcttc aattttcttt attttagaca tgggtatgaa atgtgtgtta 420  
gagttgggtt gaatgagata tacgttcaag tgaagtggca taccgttctc gagtaaggat 480  
gacctacca ttcttgagac aaatgttaca ttttagtattc agagtaaaat gtgtacctat 540

aactcaaatt cgattgacat gtatccattc aacataaaat taaaccagcc tgcacctgca 600  
tccacatttc aagtattttc aaaccgttcg gtcctatcc accgggtgta acaagacgga 660  
ttccgaattt ggaagatttt gactcaaatt cccaatttat attgaccgtg actaaatcaa 720  
ctttaacttc tataattctg attaagctcc caatttatat tcccaacggc actacctcca 780  
aaatttatag actctcatcc ctttttaaac caacttagta aacgtttttt tttttaattt 840  
tatgaagtta agtttttacc ttgtttttaa aaagaatcgt tcataagatg ccatgccaga 900  
acattagcta cacgttacac atagcatgca gccgcggaga attgtttttc ttcgccactt 960  
gtcactccct tcaaacacct aagagcttct ctctcacagc acacacatac aatcacatgc 1020  
gtgcatgcat tattacacgt gatcgccatg caaatctcct ttatagccta taaattaact 1080  
catccgcttc actctttact caaaccaaaa ctcatcaata caaacaagat taaaaacata 1140  
cacgaggatc ctcagtcaca caaagagtaa agaagaaca atg gct tcc tct atg 1194  
Met Ala Ser Ser Met  
1 5  
ctc tct tcc gct act atg gtt gcc tct ccg gct cag gcc act atg gtc 1242  
Leu Ser Ser Ala Thr Met Val Ala Ser Pro Ala Gln Ala Thr Met Val  
10 15 20  
gct cct ttc aac gga ctt aag tcc tcc gct gcc ttc cca gcc acc cgc 1290  
Ala Pro Phe Asn Gly Leu Lys Ser Ser Ala Ala Phe Pro Ala Thr Arg  
25 30 35  
aag gct aac aac gac att act tcc atc aca agc aac ggc gga cgc gtt 1338  
Lys Ala Asn Asn Asp Ile Thr Ser Ile Thr Ser Asn Gly Gly Arg Val  
40 45 50  
aac tgc atg tct aga atg cca tcc gag tcg tca gac gca gct cgt cct 1386  
Asn Cys Met Ser Arg Met Pro Ser Glu Ser Ser Asp Ala Ala Arg Pro  
55 60 65  
gcg cta aag cac gcc tac aaa cct cca gca tct gac gcc aag ggc atc 1434  
Ala Leu Lys His Ala Tyr Lys Pro Pro Ala Ser Asp Ala Lys Gly Ile  
70 75 80 85  
acg atg gcg ctg acc atc att ggc acc tgg acc gca gtg ttt tta cac 1482  
Thr Met Ala Leu Thr Ile Ile Gly Thr Trp Thr Ala Val Phe Leu His  
90 95 100  
gca ata ttt caa atc agg cta ccg aca tcc atg gac cag ctt cac tgg 1530  
Ala Ile Phe Gln Ile Arg Leu Pro Thr Ser Met Asp Gln Leu His Trp  
105 110 115  
ttg cct gtg tcc gaa gcc aca gcc cag ctt ttg ggc gga agc agc agc 1578  
Leu Pro Val Ser Glu Ala Thr Ala Gln Leu Leu Gly Gly Ser Ser Ser  
120 125 130  
cta ctg cac atc gct gca gtc ttc att gta ctt gag ttc ctg tac act 1626  
Leu Leu His Ile Ala Ala Val Phe Ile Val Leu Glu Phe Leu Tyr Thr  
135 140 145

ggt cta ttc atc acc aca cat gac gca atg cat ggc acc ata gct ttg 1674  
 Gly Leu Phe Ile Thr Thr His Asp Ala Met His Gly Thr Ile Ala Leu  
 150 155 160 165  
 agg cac agg cag ctc aat gat ctc ctt ggc aac atc tgc ata tca ctg 1722  
 Arg His Arg Gln Leu Asn Asp Leu Leu Gly Asn Ile Cys Ile Ser Leu  
 170 175 180  
 tac gcc tgg ttt gac tac agc atg ctg cat cgc aag cac tgg gag cac 1770  
 Tyr Ala Trp Phe Asp Tyr Ser Met Leu His Arg Lys His Trp Glu His  
 185 190 195  
 cac aac cat act ggc gaa gtg ggg aaa gac cct gac ttc cac aag gga 1818  
 His Asn His Thr Gly Glu Val Gly Lys Asp Pro Asp Phe His Lys Gly  
 200 205 210  
 aat ccc ggc ctt gtc ccc tgg ttc gcc agc ttc atg tcc agc tac atg 1866  
 Asn Pro Gly Leu Val Pro Trp Phe Ala Ser Phe Met Ser Ser Tyr Met  
 215 220 225  
 tcc ctg tgg cag ttt gcc cgg ctg gca tgg tgg gca gtg gtg atg caa 1914  
 Ser Leu Trp Gln Phe Ala Arg Leu Ala Trp Trp Ala Val Val Met Gln  
 230 235 240 245  
 atg ctg ggg gcg ccc atg gca aat ctc cta gtc ttc atg gct gca gcc 1962  
 Met Leu Gly Ala Pro Met Ala Asn Leu Leu Val Phe Met Ala Ala Ala  
 250 255 260  
 cca atc ttg tca gca ttc cgc ctc ttc tac ttc ggc act tac ctg cca 2010  
 Pro Ile Leu Ser Ala Phe Arg Leu Phe Tyr Phe Gly Thr Tyr Leu Pro  
 265 270 275  
 cac aag cct gag cca ggc cct gca gca ggc tct cag gtg atg gcc tgg 2058  
 His Lys Pro Glu Pro Gly Pro Ala Ala Gly Ser Gln Val Met Ala Trp  
 280 285 290  
 ttc agg gcc aag aca agt gag gca tct gat gtg atg agt ttc ctg aca 2106  
 Phe Arg Ala Lys Thr Ser Glu Ala Ser Asp Val Met Ser Phe Leu Thr  
 295 300 305  
 tgc tac cac ttt gac ctg cac tgg gag cac cac aga tgg ccc ttt gcc 2154  
 Cys Tyr His Phe Asp Leu His Trp Glu His His Arg Trp Pro Phe Ala  
 310 315 320 325  
 ccc tgg tgg cag ctg ccc cac tgc cgc cgc ctg tcc ggg cgt ggc ctg 2202  
 Pro Trp Trp Gln Leu Pro His Cys Arg Arg Leu Ser Gly Arg Gly Leu  
 330 335 340  
 gtg cct gcc ttg gca tgacctgggc cctccgctgg tgaccacagcg tctgcacaag 2257  
 Val Pro Ala Leu Ala  
 345  
 agtgtcatgg agctcgaatt tccccgatcg ttcaaacatt tggcaataaa gtttcttaag 2317  
 attgaatcct gttgccggtc ttgcgatgat tatcatataa tttctgttga attacgttaa 2377  
 gcatgtaata attaacatgt aatgcatgac gttattttatg agatggggtt ttatgattag 2437  
 agtccccgcaa ttatacatatt aatacgcgat agaaaacaaa atatagcgcg caaactagga 2497  
 taaattatcg cgcgcgggtgt catctatggt actagatcgg gaattc 2543

<210> 2  
 <211> 346  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: deduced fusion protein of  
 transit peptide + peptide with beta-carotene C-4 oxygenase activity

<400> 2

```

Met Ala Ser Ser Met Leu Ser Ser Ala Thr Met Val Ala Ser Pro Ala
 1             5             10             15

Gln Ala Thr Met Val Ala Pro Phe Asn Gly Leu Lys Ser Ser Ala Ala
      20             25             30

Phe Pro Ala Thr Arg Lys Ala Asn Asn Asp Ile Thr Ser Ile Thr Ser
      35             40             45

Asn Gly Gly Arg Val Asn Cys Met Ser Arg Met Pro Ser Glu Ser Ser
      50             55             60

Asp Ala Ala Arg Pro Ala Leu Lys His Ala Tyr Lys Pro Pro Ala Ser
      65             70             75             80

Asp Ala Lys Gly Ile Thr Met Ala Leu Thr Ile Ile Gly Thr Trp Thr
      85             90             95

Ala Val Phe Leu His Ala Ile Phe Gln Ile Arg Leu Pro Thr Ser Met
      100            105            110

Asp Gln Leu His Trp Leu Pro Val Ser Glu Ala Thr Ala Gln Leu Leu
      115            120            125

Gly Gly Ser Ser Ser Leu Leu His Ile Ala Ala Val Phe Ile Val Leu
      130            135            140

Glu Phe Leu Tyr Thr Gly Leu Phe Ile Thr Thr His Asp Ala Met His
      145            150            155            160

Gly Thr Ile Ala Leu Arg His Arg Gln Leu Asn Asp Leu Leu Gly Asn
      165            170            175

Ile Cys Ile Ser Leu Tyr Ala Trp Phe Asp Tyr Ser Met Leu His Arg
      180            185            190

Lys His Trp Glu His His Asn His Thr Gly Glu Val Gly Lys Asp Pro
      195            200            205

Asp Phe His Lys Gly Asn Pro Gly Leu Val Pro Trp Phe Ala Ser Phe
      210            215            220

Met Ser Ser Tyr Met Ser Leu Trp Gln Phe Ala Arg Leu Ala Trp Trp
      225            230            235            240

Ala Val Val Met Gln Met Leu Gly Ala Pro Met Ala Asn Leu Leu Val
      245            250            255
  
```

Ser Gly Arg Gly Leu Val Pro Ala Leu Ala  
340 345

29295/BN

# Claims

1. A DNA construct comprising in the 5' to 3' direction of transcription operably linked a promoter region directing transcription to the seed of an oilseed plant, a nucleotide sequence coding for at least one peptide with enzyme activity necessary for keto group containing xanthophyll production and esterification in an oilseed plant and a transcriptional termination region.

2. The DNA construct according to claim 1, which between the promoter region and the nucleotide sequence coding for at least one peptide with enzyme activity additionally comprises a nucleotide sequence coding for a transit peptide directing the translated fusion polypeptide to the chloroplast of the oilseed plant.

3. The DNA construct according to claim 1 or 2, wherein the promoter is a napin promoter, the peptide with enzyme activity necessary for keto group containing xanthophyll production and esterification is selected from the group consisting of peptides with, 1-D-deoxyxylulose 5-phosphate synthase, isopentenyl pyrophosphate:dimethylallyl pyrophosphate isomerase, geranylgeranyl pyrophosphate synthase, phytoene synthase, phytoene desaturase, zeta-carotene desaturase, lycopene beta-cyclase,  $\beta$ -carotene hydroxylase,  $\beta$ -carotene C-4-oxygenase, and acyl transferase activity.

4. The DNA construct according to any one of claims 1 - 3, wherein the nucleotide sequence coding for a peptide with enzyme activity is a nucleotide sequence coding for a N-terminally truncated  $\beta$ -carotene C-4-oxygenase gene from the alga *Haematococcus pluvialis*.

5. The DNA construct according to claim 4, wherein the nucleotide sequence is SEQ ID NO:1.

6. Transgenic oilseed plant cell comprising the DNA construct of any one of claims 1-5 .

7. Transgenic oilseed plant cell according to claim 6, wherein the oilseed plant is selected from the group consisting of rape, sunflower, soybean and mustard.

8. Transgenic oilseed plant-produced xanthophyll.

9. Transgenic oilseed plant-produced canthaxanthin

10. Transgenic oilseed plant-produced astaxanthin.

11. Transgenic oilseed plant-produced astaxanthin esters.

Napin promoter

AAGCTTTCTTCATCGGTGATTGATTCCTTTAAAGACTTATGTTTCTTATCTTGCTTCTGA  
GGCAAGTATTTCAGTTACCAGTTACCACTTATATTCTGGACTTTCTGACTGCATCCTCATT  
TTTCCAACATTTTAAATTTCACTATTGGCTGAATGCTTCTTCTTTGAGGAAGAAACAATT  
CAGATGGCAGAAATGTATCAACCAATGCATATATACAAATGTACCTCTTGTTCTCAAAAC  
ATCTATCGGATGGTTCCATTTGCTTTGTCATCCAATTAGTGACTACTTTATATTATTAC  
TCCTCTTTATTACTATTTTCATGCGAGGTTGCCATGTACATTATATTTGTAAGGATTGAC  
GCTATTGAGCGTTTTTCTTCAATTTTCTTATTTTAGACATGGGTATGAAATGTGTGTTA  
GAGTTGGGTTGAATGAGATATACGTTCAAGTGAAGTGGCATAACCGTTCTCGAGTAAGGAT  
GACCTACCCATTCTTGAGACAAATGTTACATTTTAGTATCAGAGTAAAATGTGTACCTAT  
AACTCAAATTCGATTGACATGTATCCATTCAACATAAAATTAAACCAGCCTGCACCTGCA  
TCCACATTTCAAGTATTTTCAAACCGTTCGGCTCCTATCCACCGGGTGTAACAAGACGGA  
TTCCGAATTTGGAAGATTTTGACTCAAATCCCAATTTATATTGACCGTGACTAAATCAA  
CTTTAACTTCTATAATTCTGATTAAGCTCCCAATTTATATTCCCAACGGCACTACCTCCA  
AAATTTATAGACTCTCATCCCCTTTTAAACCAACTTAGTAAACGTTTTTTTTTTTAAATTT  
TATGAAGTTAAGTTTTTACCTTGTTTTTAAAAAGAATCGTTCATAAGATGCCATGCCAGA  
ACATTAGCTACACGTTACACATAGCATGCAGCCGCGGAGAATTGTTTTTCTTCGCCACTT  
GTCACTCCCTTCAAACACCTAAGAGCTTCTCTCTCACAGCACACACATACAATCACATGC  
GTGCATGCATTATTACACGTGATCGCCATGCAAATCTCCTTTATAGCCTATAAATTA  
ACTCATCCGCTTCACTCTTTACTCAAACCAAACTCATCAATACAAACAAGATTAAAAACATA

End                      -28 untranslated leader                      TP start  
CACGAGGATCCTCAGTCACACAAAGAGTAAAGAAGAACAATGGCTTCCTCTATGCTCTCT  
M A S S M L S

TCCGCTACTATGGTTGCCTCTCCGGCTCAGGCCACTATGGTCGCTCCTTTCAACGGACTT  
S A T M V A S P A Q A T M V A P F N G L  
AAGTCCTCCGCTGCCTTCCCAGCCACCCGCAAGGCTAACAACGACATTACTTCCATCACA  
K S S A A F P A T R K A N N D I T S I T

FIG. 1

TP End                      C-4-Oxygenase

AGCAACGGCGGACGCGTTAACTGCATGTCTAGAATGCCATCCGAGTCGTCAGACGCAGCT  
 S N G G R V N C M S R M P S E S S D A A

CGTCCTGCGCTAAAGCACGCCTACAAACCTCCAGCATCTGACGCCAAGGGCATCACGATG  
 R P A L K H A Y K P P A S D A K G I T M

GCGCTGACCATCATTGGCACCTGGACCGCAGTGTTTTTACACGCAATATTTCAAATCAGG  
 A L T I I G T W T A V F L H A I F Q I R

CTACCGACATCCATGGACAGCTTCACTGGTTGCCTGTGTCCGAAGCCACAGCCCAGCTT  
 L P T S M D Q L H W L P V S E A T A Q L

TTGGGCGGAAGCAGCAGCCTACTGCACATCGCTGCAGTCTTCATTGTACTTGAGTTCCTG  
 L G G S S S L L H I A A V F I V L E F L

TACACTGGTCTATTCATCACCACACATGACGCAATGCATGGCACCATAGCTTTGAGGCAC  
 Y T G L F I T T H D A M H G T I A L R H

AGGCAGCTCAATGATCTCCTTGGCAACATCTGCATATCACTGTACGCCTGGTTTGACTAC  
 R Q L N D L L G N I C I S L Y A W F D Y

AGCATGCTGCATCGCAAGCACTGGGAGCACCACAACCATACTGGCGAAGTGGGGAAAGAC  
 S M L H R K H W E H H N H T G E V G K D

CCTGACTTCCACAAGGGAAATCCCGGCCTTGTCCCCTGGTTCGCCAGCTTCATGTCCAGC  
 P D F H K G N P G L V P W F A S F M S S

TACATGTCCCTGTGGCAGTTTGCCCCGCTGGCATGGTGGGCAGTGGTGATGCAAATGCTG  
 Y M S L W Q F A R L A W W A V V M Q M L

GGGGCGCCCATGGCAAATCTCCTAGTCTTCATGGCTGCAGCCCCAATCTTGTGAGCATTG  
 G A P M A N L L V F M A A A P I L S A F

CGCCTCTTCTACTTCGGCACTTACCTGCCACACAAGCCTGAGCCAGGCCCTGCAGCAGGC  
 R L F Y F G T Y L P H K P E P G P A A G

TCTCAGGTGATGGCCTGGTTCAGGGCCAAGACAAGTGAGGCATCTGATGTGATGAGTTTC  
 S Q V M A W F R A K T S E A S D V M S F

CTGACATGCTACCACTTTGACCTGCACTGGGAGCACCACAGATGGCCCTTTGCCCCCTGG  
 L T C Y H F D L H W E H H R W P F A P W

C-4 oxygenase Stop

TGGCAGCTGCCCCACTGCCGCCGCTGTCCGGGCGTGGCCTGGTGCCTGCCTTGGCATGA  
 W Q L P H C R R L S G R G L V P A L A \*

FIG.1 (cont.)



**C-4 oxygenase untranslated region** **Nos term**

**CCTGGTCCCTCCGCTGGTGACCCAGCGTCTGCACAAGAGTGTTCATGGAGCTCGAATTTCC**

**CCGATCGTTCAAACATTTGGCAATAAAGTTTCTTAAGATTGAATCCTGTTGCCGGTCTTG**

**CGATGATTATCATATAATTTCTGTTGAATTACGTTAAGCATGTAATAATTAACATGTAAT**

**GCATGACGTTATTTATGAGATGGGTTTTTATGATTAGAGTCCCGCAATTATACATTTAAT**

**ACGCGATAGAAAACAAAATATAGCGCGCAAAGTAGGATAAATTATCGCGCGCGGTGTCAT**

**end**

**CTATGTTACTAGATCGGGAATTC**

**Fig.1 (cont.)**

29295/BN

**Abstract**

5 A DNA construct comprising in the 5' to 3' direction of transcription operably linked  
a promoter region directing transcription to the seed of an oilseed plant, a nucleotide sequence  
coding for at least one peptide with enzyme activity necessary for keto group containing  
xanthophyll production and esterification in an oilseed plant and a transcriptional termination  
region is disclosed. The DNA construct may additionally comprise a nucleotide sequence  
coding for a transit peptide directing the translated fusion polypeptide to the chloroplast of the  
oilseed plant. The peptide with enzyme activity is preferably a peptide with  $\beta$ -carotene C-4-  
10 oxygenase activity, e.g. from the alga *Haematococcus pluvialis*.

Comprised by the invention are also a transgenic oilseed plant cell, e.g. of rape,  
sunflower, soybean or mustard origin; transgenic oilseed plant-produced xanthophyll;  
transgenic oilseed plant-produced canthaxanthin; transgenic oilseed plant-produced  
astaxanthin; and transgenic oilseed plant-produced astaxanthin esters.

15

PL99-09-17